

Qualitative and quantitative analysis of methanol extract of *Crataeva nurvala* stem bark[#]

K. K. Aathira¹, Bibu John Kariyil^{2*}, G. Dhanusha¹, J. S. Haima¹, S. Sujith², M. Shynu³ and A. R. Nisha⁴

Department of Veterinary Pharmacology and Toxicology,
College of Veterinary and Animal Sciences, Mannuthy, Thrissur- 680 651,
Kerala Veterinary and Animal Sciences University, Kerala, India.

Received: 21.12.2020

Accepted: 20.01.2021

Published: 01.06.2021

Citation:

Abstract

Medicinal plants are precious source of bioactive compounds which possess a range of beneficial properties and they remain the major source of medicine for a large proportion of population in this world. From ancient time, Crataeva nurvala was used as a vital herb in Ayurvedic system of medicine. According to Unani system of medicine, bark of C. nurvala is used as an appetite stimulant and as an agent to decrease the secretion of bile and phlegm. In the present study, methanol extract of stem bark of C. nurvala was analysed for preliminary phytochemicals and chemical profiling of the extract was illustrated using gas chromatography and mass spectrometry (GC-MS) analysis. The phytochemical analysis revealed that the plant extract contained alkaloids, steroids and triterpenoids. Gas chromatography mass spectrometry analysis determined the presence of different compounds of biological importance. The identification and characterisation of the phytoconstituents in the extract could pave the way for the discovery of new drugs for various ailments.

Keywords: *Crataeva nurvala*, triterpenes, alkaloids, steroids

Running title: Qualitative and quantitative analysis of *Crataeva nurvala* stem bark

[#]Part of MVSc thesis submitted to Kerala Veterinary and Animal Sciences University,
Pookode, Wayanad, Kerala

1. MVSc Scholar

2. Assistant Professor

3. Assistant Professor, Department of Veterinary Biochemistry

4. Assistant Professor and Head

*Corresponding author: bibijohn@kvasu.ac.in, Ph: 9895297842

Medicinal plants are valuable source of naturally active phytochemicals. They are the naturally occurring chemical compounds found in plants which provide health benefits for humans and animals. These compounds commonly known as secondary plant metabolites have been attributed to have different biological properties providing protection against various diseases.

Crataeva nurvala Buch-Ham., commonly known as Varuna, Neermathalam, Barna Chal, belonging to the family of Capparidacea, is a moderate sized deciduous tree. A variety of medicinal properties have been reported for *C. nurvala* and its stem bark. It has been traditionally used in treating blood flow, waste elimination, breathing problems, fever, metabolic disorders, joint lubrication and wound healing (Vashist *et al.*, 2020). Mekap *et al.* (2011) determined the antiurolithiatic activity of *C. nurvala*. Root and bark are documented to be laxative, lithotriptic and was found to increase the appetite and biliary secretion (Fletcher, 1993; Malini *et al.*, 1995). The ethanol and aqueous extracts of the dried stem bark of *C. nurvala* have been found to possess significant anti-fertility effects in rats (Bhaskar *et al.*, 2009). The antidiarrhoeal activity of ethanol extracts of *C. nurvala* stem bark has been reported by Inayathulla *et al.* (2010). *Crataeva nurvala* stem bark extract exhibited antidiabetic activity against alloxan induced diabetic albino rats in the study done by Sikarwar and Patil (2010). Thus, the present study was carried out to evaluate the various phytochemical constituents present in the bark of methanol extract of *C. nurvala* which would be helpful to delineate the various biological activities shown by the stem bark.

Materials and methods

Plant collection and identification

The bark of *C. nurvala* was collected from Valluvanad, Palakkad, Kerala (Fig. 1 and 2). The collected plant material was identified and its authenticity was confirmed by Raw Material Herbarium and Museum (RHMD), NISCAIR, New Delhi, India.

Preparation of extracts

Freshly collected bark of *C. nurvala* were cleaned to remove adhering dust and then dried under shade. The dried bark was coarsely powdered using an electric pulveriser and the powder obtained was extracted using a Soxhlet apparatus with methanol at 67 °C. The methanol extract was then concentrated using a rotary vacuum evaporator under reduced pressure and temperature (40 °C). The yield of the extract was calculated using the formula: Yield value (%) = Extracts obtained/ Total amount of crude drug × 100, and kept under refrigeration in an airtight container after complete evaporation of the solvent for further use.

Qualitative phytochemical analysis

The extracts were tested for the presence of bioactive compounds using methods described by Harborne (1998). Fifty milligrams of the extract were dissolved in 3 mL of chloroform. Few drops of concentrated sulphuric acid were added and the solution was allowed to stand. Formation of red colour directed the presence of steroids.

GC-MS analysis

The active phytochemical principles of methanol extract of *C. nurvala* was analysed using GC-MS system of Centre for Analytical Instrumentation- Kerala (CAI-K), Kerala Forest Research Institute (KFRI), Peechi, Kerala. The GC-MS analysis was carried out on Gas chromatography Mass Spectrometer (Shimadzu GC-MS, Japan, QP2010SE) with a mass range of 1.5- 1000 m/z. Helium at a flow rate of 1 mL/ min was used as the carrier gas. The oven temperature was maintained at 80 °C for 4 min and then increased to 280 °C in 6 minutes. The injector temperature was 260 °C and total analysis time was 50 minutes. Aliquot of the extract (0.4 µL) was injected into the chromatographic column after obtaining a clear baseline. The interpretation of the mass spectrum of GC-MS was guided using the database of the National Institute of Standards and Technology (NIST 11) and WILEY 8. The spectrum of the unknown compounds was related with the spectrum of the known compounds. The name and molecular weight of the compounds of the tested materials were ascertained.

[Type here]

Results and discussion

Qualitative phytochemical analysis

The qualitative phytochemical screening of methanol extract of stem bark of *C. nurvala* showed the presence of steroids, triterpenoids and alkaloids (Arunima, 2011). Phytochemical screening of methanol extract of stem bark of *C. nurvala* revealed the presence of steroid and terpenoids as well as alkaloids, phenolics, flavanoids, tannins and saponins (Hade *et al.*, 2016) which supported our results. Sodipo *et al.* (2000) have reported that alkaloids have been associated with medicinal uses for centuries and one of their common biological properties is their cytotoxicity. Huang *et al.* (2016) isolated six phytosteroids and nine known triterpenoids from the leaves of *Chisocheton cumingianus* in which chisopanoids E and F exhibited potent cytotoxicities towards MCF-7 with IC₅₀ values of 3.24 ± 1.39 and 8.85 ± 4.73 μM , and were further proved to prevent the cell proliferation, mainly by inducing apoptosis. Haque *et al.* (2008) isolated two terpenoids, phragmalin triacetate and lupeol from ethyl acetate extract of stem bark of *C. nurvala* by chromatographic techniques. Jain *et al.* (2016) suggested that terpenoids were capable of inhibiting NF κ B through different mechanisms. Khatun *et al.* (2015) evaluated the antioxidant, anthelmintic, antimicrobial and phytochemical assessment of ethanolic extract of *C. nurvala* leaves and displayed the presence of alkaloids, flavonoids, reducing sugar, saponins, steroids, tannins. The above mentioned phyto constituents are described to exhibit various pharmacological activities (Table 1).

Table 1. Results of analysis of calculi in Fourier Transform Infrared Spectrometry with Attenuated Total Reflectance (FTIR-ATR)

Functional Group Assignments	Animal no.	Reported IR wavelength (cm ⁻¹)	Standard IR wavelength of pure struvite (cm ⁻¹)
H-O-H stretching vibrations of water of crystallization	A ₁	3401.94	3270
	A ₂	3391.81	
	A ₃	3500-3350	
	A ₄	3360.29	
H-O-H stretching vibrations of a	A ₁	-	2385
	A ₂	2331.52	

[Type here]

cluster of water molecules	A ₃	2346.67	
	A ₄	2321.69	
H-O-H bending modes of vibrations	A ₁	1440.19	1445
	A ₂	1434.90	
	A ₃	1434.68	
	A ₄	1441.17	
N-H symmetric stretching vibrations in NH ₄ ⁺ units		-	2935
N-H symmetric stretching vibrations	A ₁	3401.94	3270
	A ₂	3391.81	
	A ₃	3500-3350	
	A ₄	3360.29	
N-H symmetric stretching vibrations in NH ₄ ⁺ units		-	2935
N-H bending vibration	A ₁	1670.28	1666
	A ₂	1650.70	
	A ₃	1651.8	
	A ₄	1654.67	
N-H asymmetric bending vibration in NH ₄ ⁺ units Ionic phosphate	A ₁	995.16	1010
	A ₂	1000.4	
	A ₃	1000.48	
	A ₄	1000.14	

Standard values used as per Bindhu *et al.*, 2012

GC- MS analysis

The results of GC-MS analysis of methanol extract revealed the presence of twenty-one compounds. The GC-MS chromatogram of twenty-one compounds is depicted in Fig. 3. Thymine, 3-hydroxy-2,3-dihydromaltol, 5-hydroxymethylfurfural, n-methyl-3-hydroxymethyl pyrrolidine-2-one, cytidine, methyl pentofuranoside, undecane, 6,6-dideutero-5-methyl-, 2,4-ditert-butylphenol and 3-deoxy-d-mannoic lactone were the major compounds.

Balamurugan *et al.* (2019) have done the chemical profiling of methanol bark extract of *C. nurvala* using GC-MS technique. The study revealed the presence of 8 components such as lup-

[Type here]

20 (29)-en-3ol, 2-hydroxy-4methoxybenzaldehyde, methoprene, 1'-acetonaphthone, 1, 2-bis (Trimethylsilyl) benzene, pivalate, cyclotrisiloxane, limonen-6-ol and 4-hexadecen-6-yne .

The recognised major compounds in our study possess some significant biological activities for future drug development. Zhao *et al.* (2013) showed that 5-hydroxymethylfurfural (5-HMF) induced apoptosis and G0/G1 cell cycle arrest in human melanoma A375 cells. Takuli *et al.* (2020) elucidated the antioxidant and antibacterial activity of *Woodwardia unigemmata* (Makino) along with chemical characterisation which revealed the presence of 3-hydroxy-2,3-dihydromaltol in GC-MS analysis. Azizi *et al.* (2006) performed fast gas chromatography/ time of flight mass spectrometry (TOF-GCMS) which identified N-methyl-3-hydroxymethylpyrrolidin-2-one from the oil extract of *Pithecellobium jiringan* jack seeds which was found to abolish excess free radicals and counteract oxidative damage. Su *et al.* (2005) evaluated the antioxidant activity of methanol extract of *Morinda citrifolia* (Noni) fruits and the purification of its butanol soluble partition of methanol extract contained isolates like cytidine. Shaheed *et al.* (2018) identified methyl pentofuranoside, also known as alpha-d-mannofuranoside, from methanolic fenugreek seed extract and determined its antibacterial activity against *Streptococcus agalactiae*, *Escherichia coli*, *Enterococcus cloacae* and *Proteus mirabillis*. Gas chromatography mass spectroscopic analysis exhibited the presence of undecane,6,6-dideutero-5-methyl- in *Nigella sativa*, *Allium sativum*, *Propolis* and *Olea europaea* mixture which was depicted as antibacterial and antifungal agent (Bintang *et al.*, 2018). Chuah *et al.* (2015) suggested that 2,4-di-tert-butylphenol induced oxidative stress through the generation of reactive oxygen species, which cause lipid peroxidation and membrane damage in root tissues and chloroplast in leaf tissues, thus leading to increased levels of antioxidant enzymes. Shobana *et al.* (2009) in their study identified compounds such as 3-deoxy-d-mannonic lactone and thymine from two varieties of garlic (*Ophioscordon* and *sativum*) which was found to possess antibacterial activity against enteric pathogens. The aforesaid isolated compounds from the methanol extract of *C. nurvala* stem bark seemed to own the reported biological activity and further study of these

[Type here]

phytoconstituents may demonstrate the medicinal importance in future. The biological activities of other compounds have not been reported so far and more study of these phytoconstituents might validate the significant medicinal features in forthcoming.

Conclusion

The association among the phytochemical constituents with their biological activities is now being the matter of advanced thought. *Crataeva nurvala* is a deciduous medium sized tree, traditionally used in the treatment of kidney stones, urinary tract infection and prostate related disorder. The present study has revealed *C. nurvala* to be rich in various phytochemicals. The existence of these phytochemical constituents indicated that the bark of the plant could be used in a variety of ways which would be beneficial to the population. Gas chromatography mass spectroscopic analysis revealed the presence many compounds presumed to be responsible for eliciting the traditional medicinal activities of the bark of the plant.

Acknowledgement (If any)

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Arunima, S.S. 2011. Post-natal development of lymph nodes in Kuttanad ducks. M.V.Sc.thesis, Kerala Veterinary and Animal Sciences University, Pookode, 66p.
- Bhaskar, V.H., Profulla, K.M., Balakrishnan, B.R., Balakrishnan, N. and Sangameswaran, B. 2009. Evaluation of the anti-fertility activity of stem bark of *Crataeva nurvala* buch-hum. *Afr. J. Biotechnol.* **22**: 6453-6456.
- Bintang, M., Pasaribu, F.H., Safira, U.M. and Sidhartha, T. 2018. Identification of bioactive compound from *Nigella sativa*, *Allium sativum*, *propolis* and *oleaeuropaea* mixture as antibacterial and antifungal agent. *Earth Environ. Sci.* **196**: 12041p.

- Chuah, T.S., Norhafizah, M.Z. and Ismail, B.S. 2015. Evaluation of the biochemical and physiological activity of the natural compound, 2, 4-ditert-butylphenol on weeds. *Crop Pasture Sci.* **2**: 214-223.
- Fletcher, T.F. 1993. Nervous System. In: Dellmann, H.D. (ed.), Text book of Veterinary Histology. (4th Ed.). Lea and Febiger, Philadelphia, pp. 87-107.
- Balamurugan, V., Revathi, E., Kamalakkannan, J. and Sundaresan, A. 2019. Anticancer activity of methanol extract of *Crataeva nurvala* in HeLa cell Line. *Int. J. Cell Biol. Cell. Processes.* **5**: 9-22.
- Hade, S.N., Joshi, P.A., Pilley, H.H., Wadegaonkar, V.P. and Wadegaonkar, P.A. 2016. Evaluation of *Crataeva nurvala* extracts as antioxidant, antiproteolytic and cytotoxic against hepatocarcinoma and mouse melanoma cell lines. *J. Appl. Pharm. Sci.* **6**: 189-196.
- Haque, M.E., Islam, M.N., Gupta, D.D., Hossain, M., Shekhar, H.U. and Shibib, B.A. 2008. Triterpenoids from the stem bark of *Crataeva nurvala*. *Dhaka Univ. J. Pharm. Sci.* **7**: 71-74.
- Harborne, J.B. 1998. Phytochemical methods: A guide to modern techniques of plant analysis. (3rd Ed.). Chapman and Hall, London. 320p.
- Huang, S.S., Jian, K.L., Li, R.J., Kong, L.Y. and Yang, M.H. 2016. Phytosteroids and triterpenoids with potent cytotoxicities from the leaves of *Chisocheton cumingianus*. *RSC Adv.* **8**: 6320-6328.
- Inayathulla, S.W.R., Asif, K. and Mukesh, S. 2010. Ethanolic extract of *Crataeva nurvala* root bark in experimental animals. *Int. J. Pharm. Pharm. Sci.* **2**: 158-61.
- Jain, H., Dhingra, N., Narsinghani, T. and Sharma, R. 2016. Insights into the mechanism of natural terpenoids as NF- κ B inhibitors: An overview on their anticancer potential. *J. Clin. Exp. Oncol.* **38**: 158-168.
- Khatun, F., Mahfuz-E-Alam, M., Tithi, N.S., Nasrin, N. and Asaduzzaman, M. 2015. Evaluation of phytochemical, antioxidant, anthelmintic and antimicrobial properties of *Crataeva nurvala* Buch. Ham. leaves. *Int. J. Pharm. Sci. Res.* **4**: p.1422.

- Malini, M.M., Baskar, R. and Varalakshmi, P. 1995. Effect of lupeol, a pentacyclic triterpene, on urinary enzymes in hyperoxaluric rats. *Jpn. J. Med. Sci. Biol.* **1**: 211-220.
- Mekap, S.K., Mishra, S., Sahoo, S. and Panda, P.K. 2011. Antiuro lithiatic activity of *Crataeva magna* Lour. bark. *Indian J. Nat. Prod. Resour.* **1**: 28-33
- Shaheed, K.A., Alsirraj, M.A., Allaith, S.A., Noori, N.A., Obaid, M.H., Mouhsan, Z.M. and Swedan, S.S. 2018. The biological activities of seeds extracts for fenugreek and black cumin and its inhibitory influences towards some pathogens. *Iraq Med. J.* **2**: 46-50.
- Shobana, S., Vidhya, V.G. and Ramya, M. 2009. Antibacterial activity of garlic varieties (ophioscordon and sativum) on enteric pathogens. *Curr. Res. J. Biol. Sci.* **3**: 123-126.
- Sikarwar, M.S. and Patil, M.B. 2010. Antidiabetic activity of *Crateva nurvala* stem bark extracts in alloxan-induced diabetic rats. *J. Pharm. Bioallied Sci.* **1**: p.18.
- Sodipo, O.A., Akinniyi, J.A. and Ogunbameru, J.V. 2000. Studies on certain characteristics of extracts of bark of *Pausinystalia johimbe* and *Pausinystalia macroceras* (K Schum) Pierre ex Beille. *Glob. J. Pure Applied Sci.* **1**: 83-88.
- Su, B.N., Pawlus, A.D., Jung, H.A., Keller, W.J., McLaughlin, J.L. and Kinghorn, A.D. 2005. Chemical constituents of the fruits of *Morinda citrifolia* (Noni) and their antioxidant activity. *J. Nat. Prod.* **4**: 592-595.
- Takuli, P., Khulbe, K., Kumar, P., Parki, A., Syed, A. and Elgorban, A.M. 2020. Phytochemical profiling, antioxidant and antibacterial efficacy of a native Himalayan Fern: *Woodwardia unigemmata* (Makino) Nakai. *Saudi J. Biol. Sci.* **8**: 1961-1967.
- Vashist, S., Choudhary, M., Shammi Rajpal, D.S. and Budhwar, V. 2020. Varuna (Buch. Ham.): A brief review on Phytochemistry, *Crataeva nurvala* pharmacological profile and uses in various ailments. *Asian J. Pharm. Pharmacol.* **2**: 119-125.

Zhao, L., Chen, J., Su, J., Li, L., Hu, S., Li, B., Zhang, X., Xu, Z. and Chen, T. 2013. *In vitro* antioxidant and antiproliferative activities of 5-hydroxymethylfurfural. *J. Agric. Food Chem.* **44**: 10604-10611.



Fig. 1. Leaves and flower of *Crataeva nurvala* Buch-Ham.



Courtesy:Dept. of Dravyaguna,
College of Ayurved, Bharati Vidyapeeth, Pune

Fig. 2. Bark of *Crataeva nurvala* Buch-Ham.

