



Effect of dietary incorporation of ksheerabala residue on nutrient digestibility and blood biochemical profile in Malabari kids*

T.J. Roshma¹, K. Ally², Thirupathy Venkatachalapathy³, K. Shyama⁴ and K. George Sherin⁵.

Department of Animal Nutrition, College of Veterinary and Animal Sciences, Mannuthy, Thrissur, Kerala
Kerala Veterinary and Animal Sciences University

Citation: Roshma, T.J., Ally, K., Thirupathy, V., Shyama, K. and George, S. K. 2020. Effect of dietary incorporation of ksheerabala residue on nutrient digestibility and blood biochemical profile in Malabari kids. *J. Vet. Anim. Sci.* **51**(2): 196-200.

Received : 07.03.2020

Accepted : 09.04.2020

Published : 01-07-2020

Abstract

An experiment was conducted with eighteen weaned Malabari kids of three months of age for 90 days to assess the effect of dietary incorporation of Ksheerabala residue on nutrient digestibility and blood biochemical profile in goats. Kids were divided into three groups (T_1 , T_2 and T_3) as uniformly as possible with regard to age, sex and body weight and were offered kid starter containing Ksheerabala residue at 0, 10 and 20 per cent, respectively. Kid starters were made isonitrogenous and isocaloric (24 per cent CP and 70 per cent TDN). Kids were fed as per ICAR standards (Ranjhan, 1998). Green grass was offered as the sole source of roughage. Data on digestibility of nutrients and haematological parameters like haemoglobin, plasma protein, serum calcium, serum phosphorus, serum cholesterol and triglycerides were the criteria employed for evaluation and they did not show any significant difference ($P > 0.05$) among the groups. It could be inferred that digestibility of nutrients and haematological parameters of kids were not influenced by the inclusion of Ksheerabala residue in kid starter.

Key words: Blood parameters, Digestibility, kid starter and Ksheerabala residue

Kerala has various ayurvedic pharmaceuticals and byproducts composed of residues of medicinal herbs. Ksheerabala residue is a byproduct obtained during the preparation of ksheerabala oil which is made by incorporating *Sida cordifolia*, cow milk and gingelly oil. This residue is available in considerable quantity locally, around 800-1000 kg per week (Kottakkal Aryavudiasala, Kottakkal, Kerala) and many of the farmers are using for feeding goats. The feeding value of these residues as potential non conventional feed resource (NCFR) are yet to be explored. The main problem

*Forms part of the MVSc thesis of the first author submitted to Kerala Veterinary and Animal Sciences University, Pookode, Wayanad, Kerala

Correspondence E-mail: ally@kvasu.ac.in Mob:9447476508

1. Veterinary Surgeon, Animal Husbandry Department, Kerala
2. Professor and Head
3. Professor and Head, University Sheep and Goat Farm, Mannuthy
4. Professor and Head, Cattle Breeding Farm, Thumburmuzhi
5. Assistant Professor, Base Farm, Kolahalamedu

Copyright: © 2020 Roshma et al. This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

with residues from ayurvedic products are their palatability and probable cumulative toxic effects. The microflora of goats' rumen can utilize the fibrous residues in the byproducts of ayurvedic preparations and can possibly detoxify the intrinsic factors present in these residues to some extent (Abreu et al.2004) Hence the present study is planned to evaluate the effect of dietary incorporation of Ksheerabala residue as a NCFR in the diet of kids on nutrient digestibility and blood biochemical profile.

Materials and Methods

Eighteen healthy kids of three months of age around 8.25 kg body weight, selected from University Goat and Sheep Farm, College of Veterinary and Animal Sciences, Mannuthy, formed the experimental subjects for the study. Kids were weaned and housed individually in well ventilated, clean and dry shed with facilities for feeding and watering. They were divided into three groups of six animals each as uniformly as possible with regard to age, sex and body weight and were allotted randomly to one of the three groups, T₁ (kid starter), T₂ (kid starter containing 10 per cent ksheerabala residue) and T₃ (kid starter containing 20 per cent ksheerabala residue). All the rations were made isonitrogenous and isocaloric (24 per cent CP and 70 per cent TDN). Good quality green grass and clean drinking water were offered from first week of age. Ksheerabala residue available after extraction of oil from the pharmaceutical firm (Kottakkal Aryavydiasala, Kerala) was procured freshly as a firm solid block of around 35-40 kg weight. Proximate composition of Ksheerabala residue is presented in Table 1. The ingredient and chemical composition of experimental rations are presented in Table 2.

The experimental animals were maintained on the respective treatments for a period of 90 days. Daily records of concentrate and grass given and balance if any were maintained throughout the feeding trial. Data on daily dry matter intake was maintained throughout the experimental period. A digestibility trial involving five days collection period was conducted at 12th week of the study period to assess the digestibility coefficient of nutrients. Representative samples of kid starter and green grass offered were taken daily

during the digestion trial for chemical analysis. The balance of feed and grass samples were collected from individual animals and their moisture content was determined daily. At the end of the collection period feed samples collected daily were pooled and subjected to chemical analysis. The dung was collected manually as and when it was voided. All precautions were taken to collect the dung quantitatively, uncontaminated with urine, feed residue or dirt. The dung collected each day was weighed accurately and were kept in double lined air tight plastic bags and stored fresh in deep freezer during the entire collection period. At the end of collection period daily samples stored from each animal were pooled and used for chemical analysis. Kid starter, fodder and dung samples were analyzed for proximate principles (AOAC, 2012). The acid detergent fiber (ADF) was estimated by the method suggested by Van Soest (1963) and neutral detergent fiber (NDF) by the method suggested by Van Soest and Whine (1967).

Blood samples were collected from all animals at the end of the experiment. These samples were used to determine haemoglobin, plasma total protein (Jong and Vegeter, 1950), calcium, phosphorus (Bernhart and Wreath, 1955), plasma cholesterol (Lie *et al.*, 1976) and triglycerides (McGowan *et al.*, 1983) Blood haemoglobin was estimated by cyanomethaemoglobin method using reagents from Agappe diagnostics Ltd, Ernakulam, India. Plasma protein, calcium, phosphorus, serum cholesterol and serum triglycerides were determined using the blood analyzer (Mispa plus, SEAC radim group) and kits supplied by same suppliers. Data gathered on various parameters were analyzed statistically using Analysis of Variance (Snedecor and Cochran, 1994).

Results and Discussion

From the data collected on daily dry matter intake, the weekly average daily dry matter intake of the kids were 0.24, 0.24 and 0.22 kg, respectively in T₁, T₂ and T₃ initially and the corresponding values at the end of feeding trial were 0.56, 0.48 and 0.52 kg, which were similar.

Table 1. Proximate composition of Ksheerabala residue (%on dry matter basis)

Dry matter	92.55
Crude protein	29.52
Ether extract	13.26
Crude fibre	6.39
Total ash	8.42
Nitrogen Free Extract	42.41
Acid insoluble ash	0.06

Nutrient digestibility

Digestibility of dry matter, crude protein, crude fibre, ether extract, nitrogen free extract, NDF and ADF observed in the present study were 77.34, 76.43 and 75.28, 81.97, 81.36 and 79.66, 64.47, 64.29 and 61.61, 81.33, 82.78 and 84.65, 84.04, 82.58 and 81.65, 59.77, 58.20 and 57.86, 55.58, 55.23 and 52.78 per cent in T₁, T₂ and T₃ kids, respectively and statistical analysis did not reveal any significant difference

between the groups (Table 3). In agreement with the present result, Rani *et al.* (2006) reported that digestibility of DM, CP, NDF and ADF were similar in buffalo calves supplemented with two herbs namely bringraj (*Eclipta alba*) and kutki (*Kutki picorrhiza*) at the rate of 0.4 per cent of dry matter intake. Verma *et al.* (1995) who also found that inclusion of water washed neem seed kernel cake at 0, 15 and 25 per cent levels in goat ration had similar effect on digestibility of DM, CP, EE and NFE.

Obeidat and Gharaybeh (2011) observed that the digestibility coefficient of DM and CP were similar in kids fed with diet containing 0, 10 and 20 per cent of sesame hull whereas the EE digestibility was significantly higher in treatment groups compared to control. Tufarelli *et al.* (2012) also reported that digestibility of CP and EE were similar in lambs fed with partly destoned exhausted olive cake

Table 2. Ingredient and chemical composition of kid starters

Ingredients	Percentage composition of calf starter		
	T ₁	T ₂	T ₃
Maize	35	29	23
Wheat bran	25	26	28
Soya bean meal	28	23	17
Dried fish	9	9	9
Ksheerabala residue	0	10	20
Salt	1	1	1
Calcite	2	2	2
Total	100.00	100.00	100.00
Vitamin Supplement g/100 kg feed	25	25	25
Trace mineral mix g/100 kg feed	25	25	25
Toxin binder g/100 kg feed	100	100	100
Salinomycin sodium g/100 kg feed	50	50	50
Chemical composition*			
Dry matter	92.19	91.73	91.93
Crude protein	23.74	23.99	24.51
Ether extract	4.95	5.38	6.04
Crude fibre	5.43	5.71	6.08
Total ash	9.72	10.07	10.41
Nitrogen free extract	55.65	54.84	52.87
Acid insoluble ash	1.11	1.23	1.28
Neutral detergent fibre	22.34	22.83	25.84
Acid detergent fibre	6.61	7.81	9.48
Calcium	1.03	1.12	1.18
Phosphorus	0.65	0.70	0.86

*On dry matter basis

Table 3. Digestibility coefficient of nutrients in experimental ration*

Item	Dietary treatments		
	T ₁	T ₂	T ₃
DM	77.34 ± 0.21	76.43 ± 1.11	75.28 ± 1.26
CP	81.97 ± 0.85	81.36 ± 1.57	79.66 ± 1.03
CF	64.47 ± 0.84	64.29 ± 2.55	61.61 ± 1.56
EE	81.33 ± 0.97	82.78 ± 1.23	84.65 ± 0.94
NFE	84.04 ± 0.26	82.58 ± 1.19	81.65 ± 1.83
NDF	59.77±0.61	58.20±3.01	57.86±1.57
ADF	55.58±1.05	55.23±2.67	52.78±1.64

*on dry matter basis

Table 4. Haematological parameters of experimental kids

Parameters	T ₁	T ₂	T ₃
Haemoglobin, g/dl	11.88 ± 0.31	12.00 ± 0.46	11.94 ± 0.57
plasma protein, g/dl	6.38 ± 0.19	6.33 ± 0.2	6.28 ± 0.25
Calcium, mg/dl	10.30 ± 0.11	10.31 ± 0.15	10.24 ± 0.32
Phosphorus, mg/dl	6.02 ± 0.12	6.02 ± 0.13	5.96 ± 0.29
Cholesterol, mg/dl	110.83 ± 2.96	104.03 ± 2.93	108.62 ± 6.23
Triglyceride, mg/dl	40.69 ± 0.41	39.36 ± 0.9	40.53 ± 1.3

at 0, 10 and 15 per cent levels in ration whereas the digestibility of DM was significantly lower in both diets containing olive cake compared to control.

Haematological parameters

Data on haematological studies were documented in Table 4. The average blood haemoglobin and plasma protein concentration at the end of the experiment for group T₁, T₂ and T₃ were 11.88, 12.00 and 11.94, 6.38, 6.33 and 6.28g/ dl, respectively. The average serum calcium values in experimental animals were 10.30, 10.31 and 10.24 mg/dl for group T₁, T₂ and T₃, respectively. The average serum phosphorus concentrations in the experimental kids were 6.02, 6.02 and 5.96 mg/dl for group T₁, T₂ and T₃, respectively. The average serum cholesterol and triglyceride values recorded at the end of the experiment were 110.83, 104.03 and 108.62, 40.69, 39.36 and 40.53 mg/ dl, respectively for group T₁, T₂ and T₃. There was no significant difference (P >0.05) in any of the haematological parameters between the three groups.

Verma *et al.* (1995) studied the effect of dietary inclusion of water washed neem seed kernel

cake in growing goat ration at 0, 15 and 25 per cent levels and found that haemoglobin and cholesterol concentration were similar in treatment and control groups which is in accordance with the present result but total protein concentration was significantly lower in goats fed with neem seed kernel cake. An *et al.* (2001) found that the total cholesterol level was significantly reduced (113.92 mg/ dl) in broilers fed with ration containing 10 per cent red pepper seed oil meal compared to control diet (137.5 mg/ dl). Jasmine *et al.* (2018) reported similar values for haemoglobin, total protein, serum calcium and phosphorus in growing kids fed with ration containing spent rosemary (*Rosmarinus officianalis.L*) leaves

Conclusion

Critical evaluation of the results obtained in the present study revealed that inclusion of Ksheerabala residue in kid starter had similar effect on nutrient digestibility and haematological parameters. On summarizing the overall results of the study, it could be inferred that Ksheerabala residue can be included in the goat ration up to 20 per cent level without any adverse effect on nutrient digestibility and blood biochemical profile.

References

- Abreu. A . Carulla, J.E. Lascano, C.E.. Diaz, T. E.. Kreuzer, M. and Hess, H. D. (2004). Effects of *Sapinduss aponaria* fruits on ruminal fermentation and duodenal nitrogen flow of sheep fed a tropical grass diet with and without legume. *J. Anim. Sci.* 82:1392-1400
- An, B. K., Im, H. J. and Kang, C.W. 2001). Nutritional values of red pepper seed oil meal and effects of its supplementation on performance and physiological responses of broiler chicks. *Asian-Aust. J. Anim. Sci.* 20(6): 971-975.
- AOAC. 2012. *Official Methods of Analysis*, (20th Ed.). Association of official analytical chemists. Washington, D.C, 587p.
- Bernhart, D.N. and Wreath, A. 1955. Colorimetric determination of phosphorus by modified phosphomolybdate method. *Analyt. Chem.* 27: 440-441.
- Jasmine, M.S.S., Shyama K., Gangadevi P., Ally K and Thirupathy Venkatachalapathy R. 2018. Effect of incorporation of spent rose mary leaves on serum biochemical parameters of Malabary kids. . *J.Vet. Anim.Sci.* 49 (7):67-70
- Jong, H.H. and Vegeter, J.J. 1950. Determination of serum proteins with the biuret test . *Pharm. weekbl.* 85: 755-764.
- Lie, R.F., Schmitz, J.M., Pierre, K.J. and Gochman, N. 1976. Cholesterol oxidase based determination, by continuous-flow analysis, of total and free cholesterol in serum. *Clin. Chem.* 22: 1627-1630.
- McGowan, M.W., Artiss, J.D., Strandbergh, D.R. and Zak, B. 1983. A peroxidase coupled method for the colorimetric determination of serum triglycerides. *Clin. Chem.* 29: 538-542.
- Obeidat, B.S. and Gharaybeh, F.F. 2011. Effect of feeding sesame hull on growth performance, nutrient digestibility, and carcass characteristics of black goat kids. *Asian-Aust. J. Anim. Sci.* 24(2): 206-213.
- Rani, N., Wadhwa, M., Kaushal, S. and Bakshi, M.P.S. 2006. Herbal feed additives and growth of buffalo calves. *Anim. Nutr. Feed Tech.* 6 : 147-151.
- Ranjhan, S.K. 1998. *Nutrient Requirements of Livestock and Poultry*. (2nd Ed.). Indian Council of Agricultural Research, New Delhi. 72p.
- Snedecor, G.W. and Cochran, W.G. 1994. *Statistical Methods*. (8th Ed.). The Iowa State University press, Ames, IA. 314p.
- Tufarelli, V., Introna, M., Cazzato, E., Mazzei, D. and Laudadio, V. 2012. Suitability of partly destined exhausted olive cake by product feed ingredient for lamb production. *J. Anim.Sci.* 91: 872-877.
- Van Soest, P.J. 1963. Use of detergents in analysis of fibrous feeds. II. A rapid method for the determination of fibre and lignin. *J. Assoc. Off. Analyt. Chem.* 46: 829p
- Van Soest, P.J. and Whine, R.H. 1967. Use of detergents in the analysis of fibrous feeds. IV. The determination of plant cell wall constituents. *J. Assoc. Off. Analyt. Chem.* 50: 50p.
- Verma, A. K., Sastry, V.R.B. and Agarwal, D. K. 1995. Feeding of water washed neem (*Azadiracta indica*) seed kernel cake to growing goats. *J. Small Rum. Res.* 15: 105-11. ■